


Set 9:

CMB and Large Scale Structure


CMB Temperature Anisotropy

- Planck 2015 map of the temperature anisotropy (first discovered by COBE) from recombination:



CMB Temperature Anisotropy

- Power spectrum shows characteristic scales where the intensity of variations peak - reveals geometry and contents of the universe:




CMB Parameter Inferences

- Spectrum constrains the matter-energy contents of the universe

Parameter	[1] <i>Planck</i> TT+lowP	[2] <i>Planck</i> TE+lowP	[3] <i>Planck</i> EE+lowP	[4] <i>Planck</i> TT,TE,EE+lowP
$\Omega_b h^2$	0.02222 ± 0.00023	0.02228 ± 0.00025	0.0240 ± 0.0013	0.02225 ± 0.00016
$\Omega_c h^2$	0.1197 ± 0.0022	0.1187 ± 0.0021	$0.1150^{+0.0048}_{-0.0055}$	0.1198 ± 0.0015
$100\theta_{MC}$	1.04085 ± 0.00047	1.04094 ± 0.00051	1.03988 ± 0.00094	1.04077 ± 0.00032
τ	0.078 ± 0.019	0.053 ± 0.019	$0.059^{+0.022}_{-0.019}$	0.079 ± 0.017
$\ln(10^{10} A_s)$	3.089 ± 0.036	3.031 ± 0.041	$3.066^{+0.046}_{-0.041}$	3.094 ± 0.034
n_s	0.9655 ± 0.0062	0.965 ± 0.012	0.973 ± 0.016	0.9645 ± 0.0049
H_0	67.31 ± 0.96	67.73 ± 0.92	70.2 ± 3.0	67.27 ± 0.66
Ω_m	0.315 ± 0.013	0.300 ± 0.012	$0.286^{+0.027}_{-0.038}$	0.3156 ± 0.0091
σ_8	0.829 ± 0.014	0.802 ± 0.018	0.796 ± 0.024	0.831 ± 0.013
$10^9 A_s e^{-2\tau}$	1.880 ± 0.014	1.865 ± 0.019	1.907 ± 0.027	1.882 ± 0.012


Galaxy Redshift Surveys

- Galaxy redshift surveys (e.g. 2dF and SDSS) measure the three dimensional distribution of galaxies today:



Galaxy Power Spectrum

- SDSS LRG and Main power spectrum:



Structure Formation

- Small perturbations from inflation over the course of the 14Gyr life of the universe are gravitationally enhanced into all of the structure seen today
- Cosmic microwave background shows a snapshot at a few hundred thousand years old at recombination
- Discovery in 1992 of cosmic microwave background anisotropy provided the observational breakthrough - convincing support for adiabatic initial density fluctuations of amplitude 10^{-5}
- Combine with galaxy clustering - large scale structure seen in galaxy surveys - right amplitude given cold dark matter
- Following notes are at a slightly more advanced level than the book and are provided here for completeness

Angular Power Spectrum

- Angular distribution of radiation is essentially the 3D temperature field projected onto a shell at the distance from the observer to recombination: called the last scattering surface
- Take the radiation distribution at last scattering to also be described by an isotropic temperature fluctuation field $\Theta(\mathbf{x})$ and recombination to be instantaneous

$$\Theta(\hat{\mathbf{n}}) = \int dD \Theta(\mathbf{x}) \delta(D - D_*)$$

where D is the comoving distance and D_* denotes recombination.

- Describe the temperature field by its Fourier moments

$$\Theta(\mathbf{x}) = \int \frac{d^3k}{(2\pi)^3} \Theta(\mathbf{k}) e^{i\mathbf{k}\cdot\mathbf{x}}$$

Angular Power Spectrum

- Power spectrum

$$\langle \Theta(\mathbf{k})^* \Theta(\mathbf{k}') \rangle = (2\pi)^3 \delta(\mathbf{k} - \mathbf{k}') P_T(k)$$

$$\Delta_T^2 = k^3 P_T / 2\pi^2$$

- Temperature field

$$\Theta(\hat{\mathbf{n}}) = \int \frac{d^3k}{(2\pi)^3} \Theta(\mathbf{k}) e^{i\mathbf{k} \cdot D_* \hat{\mathbf{n}}}$$

- Multipole moments $\Theta(\hat{\mathbf{n}}) = \sum_{\ell m} \Theta_{\ell m} Y_{\ell m}$
- Expand out plane wave in spherical coordinates

$$e^{i\mathbf{k} D_* \cdot \hat{\mathbf{n}}} = 4\pi \sum_{\ell m} i^\ell j_\ell(k D_*) Y_{\ell m}^*(\mathbf{k}) Y_{\ell m}(\hat{\mathbf{n}})$$

Angular Power Spectrum

- Power spectrum

$$\Theta_{\ell m} = \int \frac{d^3 k}{(2\pi)^3} \Theta(\mathbf{k}) 4\pi i^\ell j_\ell(k D_*) Y_{\ell m}(\mathbf{k})$$

$$\begin{aligned} \langle \Theta_{\ell m}^* \Theta_{\ell' m'} \rangle &= \int \frac{d^3 k}{(2\pi)^3} (4\pi)^2 (i)^{\ell-\ell'} j_\ell(k D_*) j_{\ell'}(k D_*) Y_{\ell m}^*(\mathbf{k}) Y_{\ell' m'}(\mathbf{k}) P_T(k) \\ &= \delta_{\ell\ell'} \delta_{mm'} 4\pi \int d \ln k j_\ell^2(k D_*) \Delta_T^2(k) \end{aligned}$$

with $\int_0^\infty j_\ell^2(x) d \ln x = 1/(2\ell(\ell+1))$, slowly varying Δ_T^2

- Angular power spectrum:

$$C_\ell = \frac{4\pi \Delta_T^2(\ell/D_*)}{2\ell(\ell+1)} = \frac{2\pi}{\ell(\ell+1)} \Delta_T^2(\ell/D_*)$$

Thomson Scattering

- Thomson scattering of photons off of free electrons is the most important CMB process with a cross section (averaged over polarization states) of

$$\sigma_T = \frac{8\pi\alpha^2}{3m_e^2} = 6.65 \times 10^{-25} \text{cm}^2$$

- Density of free electrons in a fully ionized $x_e = 1$ universe

$$n_e = (1 - Y_p/2)x_e n_b \approx 10^{-5} \Omega_b h^2 (1+z)^3 \text{cm}^{-3},$$

where $Y_p \approx 0.24$ is the Helium mass fraction, creates a high (comoving) Thomson opacity

$$\dot{\tau} \equiv n_e \sigma_T a$$

where dots are conformal time $\eta \equiv \int dt/a$ derivatives and τ is the optical depth.

Tight Coupling Approximation

- Near **recombination** $z \approx 10^3$ and $\Omega_b h^2 \approx 0.02$, the (comoving) **mean free path** of a photon

$$\lambda_C \equiv \frac{1}{\dot{\tau}} \sim 2.5 \text{Mpc}$$

small by cosmological standards!

- On scales $\lambda \gg \lambda_C$ photons are **tightly coupled** to the electrons by Thomson scattering which in turn are tightly coupled to the baryons by Coulomb interactions
- Specifically, their bulk velocities are defined by a **single fluid velocity** $v_\gamma = v_b$ and the photons carry **no anisotropy** in the rest frame of the baryons
- \rightarrow No **heat conduction** or **viscosity** (anisotropic stress) in fluid

Zeroth Order Approximation

- Momentum density of a fluid is $(\rho + p)v$, where p is the pressure
- Neglect the momentum density of the baryons

$$R \equiv \frac{(\rho_b + p_b)v_b}{(\rho_\gamma + p_\gamma)v_\gamma} = \frac{\rho_b + p_b}{\rho_\gamma + p_\gamma} = \frac{3\rho_b}{4\rho_\gamma}$$
$$\approx 0.6 \left(\frac{\Omega_b h^2}{0.02} \right) \left(\frac{a}{10^{-3}} \right)$$

since $\rho_\gamma \propto T^4$ is fixed by the CMB temperature $T = 2.73(1 + z)\text{K}$
– OK substantially before recombination

- Neglect radiation in the expansion

$$\frac{\rho_m}{\rho_r} = 3.6 \left(\frac{\Omega_m h^2}{0.15} \right) \left(\frac{a}{10^{-3}} \right)$$

- Neglect gravity

Fluid Equations

- Density $\rho_\gamma \propto T^4$ so define temperature fluctuation Θ

$$\delta_\gamma = 4 \frac{\delta T}{T} \equiv 4\Theta$$

- Real space continuity eqn. transformed to Fourier space $\nabla \rightarrow i\mathbf{k}$

$$\dot{\delta}_\gamma = -(1 + w_\gamma)k v_\gamma$$

$$\dot{\Theta} = -\frac{1}{3}k v_\gamma$$

- Euler equation (neglecting gravity)

$$\dot{v}_\gamma = -(1 - 3w_\gamma)\frac{\dot{a}}{a}v + \frac{kc_s^2}{1 + w_\gamma}\delta_\gamma$$

$$\dot{v}_\gamma = kc_s^2 \frac{3}{4}\delta_\gamma = 3c_s^2 k\Theta$$

Oscillator: Take One

- Combine these to form the simple harmonic oscillator equation

$$\ddot{\Theta} + c_s^2 k^2 \Theta = 0$$

where the sound speed is adiabatic

$$c_s^2 = \frac{\delta p}{\delta \rho} = \frac{\dot{p}_\gamma}{\dot{\rho}_\gamma}$$

here $c_s^2 = 1/3$ since we are photon-dominated

- General solution:

$$\Theta(\eta) = \Theta(0) \cos(k s) + \frac{\dot{\Theta}(0)}{k c_s} \sin(k s)$$

where the sound horizon is defined as $s \equiv \int c_s d\eta$

Harmonic Extrema

- All modes are **frozen** in at recombination (denoted with a subscript *) yielding temperature perturbations of **different amplitude** for different modes. For the adiabatic (curvature mode) $\dot{\Theta}(0) = 0$

$$\Theta(\eta_*) = \Theta(0) \cos(k s_*)$$

- Modes caught in the **extrema** of their oscillation will have enhanced fluctuations

$$k_n s_* = n\pi$$

yielding a **fundamental scale** or frequency, related to the inverse **sound horizon**

$$k_A = \pi / s_*$$

and a **harmonic relationship** to the other extrema as 1 : 2 : 3...

Peak Location

- The fundamental **physical scale** is translated into a fundamental **angular scale** by simple projection according to the angular diameter distance D_A

$$\theta_A = \lambda_A / D_A$$

$$\ell_A = k_A D_A$$

- In a flat universe, the distance is simply $D_A = D \equiv \eta_0 - \eta_* \approx \eta_0$, the horizon distance, and $k_A = \pi / s_* = \sqrt{3}\pi / \eta_*$ so

$$\theta_A \approx \frac{\eta_*}{\eta_0}$$

- In a **matter-dominated** universe $\eta \propto a^{1/2}$ so $\theta_A \approx 1/30 \approx 2^\circ$ or

$$\ell_A \approx 200$$

Curvature

- In a **curved universe**, the apparent or **angular diameter distance** is no longer the conformal distance $D_A = R \sin(D/R) \neq D$
- Objects in a **closed universe** are **further** than they appear! gravitational **lensing** of the background...
- Curvature scale of the universe must be substantially **larger than current horizon**
- **Flat universe** indicates critical density and implies missing energy given local measures of the matter density “**dark energy**”
- D also depends on **dark energy density** Ω_{DE} and **equation of state** $w = p_{\text{DE}}/\rho_{\text{DE}}$.
- Expansion rate at recombination or **matter-radiation ratio** enters into calculation of k_A .

Doppler Effect

- Bulk motion of fluid changes the observed temperature via Doppler shifts

$$\left(\frac{\Delta T}{T}\right)_{\text{dop}} = \hat{\mathbf{n}} \cdot \mathbf{v}_\gamma$$

- Averaged over directions

$$\left(\frac{\Delta T}{T}\right)_{\text{rms}} = \frac{v_\gamma}{\sqrt{3}}$$

- Acoustic solution

$$\begin{aligned} \frac{v_\gamma}{\sqrt{3}} &= -\frac{\sqrt{3}}{k} \dot{\Theta} = \frac{\sqrt{3}}{k} k c_s \Theta(0) \sin(ks) \\ &= \Theta(0) \sin(ks) \end{aligned}$$

Doppler Peaks?

- Doppler effect for the photon dominated system is of equal amplitude and $\pi/2$ out of phase: extrema of temperature are turning points of velocity
- Effects add in quadrature:

$$\left(\frac{\Delta T}{T}\right)^2 = \Theta^2(0)[\cos^2(ks) + \sin^2(ks)] = \Theta^2(0)$$

- No peaks in k spectrum! However the Doppler effect carries an angular dependence that changes its projection on the sky

$$\hat{\mathbf{n}} \cdot \mathbf{v}_\gamma \propto \hat{\mathbf{n}} \cdot \hat{\mathbf{k}}$$

- Coordinates where $\hat{\mathbf{z}} \parallel \hat{\mathbf{k}}$

$$Y_{10}Y_{\ell 0} \rightarrow Y_{\ell \pm 1 0}$$

recoupling $j'_\ell Y_{\ell 0}$: no peaks in Doppler effect

Restoring Gravity

- Take a simple photon dominated system with gravity
- Continuity altered since a gravitational potential represents a stretching of the spatial fabric that dilutes number densities – formally a spatial curvature perturbation
- Think of this as a perturbation to the scale factor $a \rightarrow a(1 + \Phi)$ so that the cosmological redshift is generalized to

$$\frac{\dot{a}}{a} \rightarrow \frac{\dot{a}}{a} + \dot{\Phi}$$

so that the continuity equation becomes

$$\dot{\Theta} = -\frac{1}{3}k v_{\gamma} - \dot{\Phi}$$

Restoring Gravity

- Gravitational force in momentum conservation $\mathbf{F} = -m\nabla\Psi$ generalized to momentum density modifies the Euler equation to

$$\dot{v}_\gamma = k(\Theta + \Psi)$$

- General relativity says that Φ and Ψ are the relativistic analogues of the Newtonian potential and that $\Phi \approx -\Psi$.
- In our matter-dominated approximation, Φ represents matter density fluctuations through the cosmological Poisson equation

$$k^2\Phi = 4\pi G a^2 \rho_m \Delta_m$$

where the difference comes from the use of comoving coordinates for k (a^2 factor), the removal of the background density into the background expansion ($\rho\Delta_m$) and finally a coordinate subtlety that enters into the definition of Δ_m

Constant Potentials

- In the matter dominated epoch potentials are constant because infall generates velocities as $v_m \sim k\eta\Psi$
- Velocity divergence generates density perturbations as $\Delta_m \sim -k\eta v_m \sim -(k\eta)^2\Psi$
- And density perturbations generate potential fluctuations as $\Phi \sim \Delta_m/(k\eta)^2 \sim -\Psi$, keeping them constant. Note that because of the expansion, density perturbations must grow to keep potentials constant.
- Here we have used the Friedman equation $H^2 = 8\pi G\rho_m/3$ and $\eta = \int d\ln a/(aH) \sim 1/(aH)$
- More generally, if stress perturbations are negligible compared with density perturbations ($\delta p \ll \delta\rho$) then potential will remain roughly constant – more specifically a variant called the Bardeen or comoving curvature \mathcal{R} is constant

Oscillator: Take Two

- Combine these to form the **simple harmonic oscillator** equation

$$\ddot{\Theta} + c_s^2 k^2 \Theta = -\frac{k^2}{3} \Psi - \ddot{\Phi}$$

- In a **CDM dominated** expansion $\dot{\Phi} = \dot{\Psi} = 0$. Also for **photon domination** $c_s^2 = 1/3$ so the oscillator equation becomes

$$\ddot{\Theta} + \ddot{\Psi} + c_s^2 k^2 (\Theta + \Psi) = 0$$

- Solution is just an **offset version** of the original

$$[\Theta + \Psi](\eta) = [\Theta + \Psi](0) \cos(ks)$$

- $\Theta + \Psi$ is also the **observed temperature fluctuation** since photons lose energy climbing out of **gravitational potentials** at recombination

Effective Temperature

- Photons climb out of potential wells at last scattering
- Lose energy to gravitational redshifts
- Observed or effective temperature

$$\Theta + \Psi$$

- Effective temperature oscillates around zero with amplitude given by the initial conditions
- Note: initial conditions are set when the perturbation is outside of horizon, need inflation or other modification to matter-radiation FRW universe.
- GR says that initial temperature is given by initial potential

Sachs-Wolfe Effect and the Magic 1/3

- A gravitational potential is a perturbation to the temporal coordinate [formally a gauge transformation]

$$\frac{\delta t}{t} = \Psi$$

- Convert this to a perturbation in the scale factor,

$$t = \int \frac{da}{aH} \propto \int \frac{da}{a\rho^{1/2}} \propto a^{3(1+w)/2}$$

where $w \equiv p/\rho$ so that during matter domination

$$\frac{\delta a}{a} = \frac{2}{3} \frac{\delta t}{t}$$

- CMB temperature is cooling as $T \propto a^{-1}$ so

$$\Theta + \Psi \equiv \frac{\delta T}{T} + \Psi = -\frac{\delta a}{a} + \Psi = \frac{1}{3}\Psi$$

Baryon Loading

- Baryons add extra **mass** to the photon-baryon fluid
- Controlling parameter is the **momentum density ratio**:

$$R \equiv \frac{p_b + \rho_b}{p_\gamma + \rho_\gamma} \approx 30\Omega_b h^2 \left(\frac{a}{10^{-3}} \right)$$

of order **unity** at recombination

- Momentum density of the **joint system** is conserved

$$\begin{aligned} (\rho_\gamma + p_\gamma)v_\gamma + (\rho_b + p_b)v_b &\approx (p_\gamma + p_\gamma + \rho_b + \rho_\gamma)v_\gamma \\ &= (1 + R)(\rho_\gamma + p_\gamma)v_{\gamma b} \end{aligned}$$

where the controlling parameter is the **momentum density ratio**:

$$R \equiv \frac{p_b + \rho_b}{p_\gamma + \rho_\gamma} \approx 30\Omega_b h^2 \left(\frac{a}{10^{-3}} \right)$$

of order **unity** at recombination

New Euler Equation

- Momentum density ratio enters as

$$[(1 + R)v_{\gamma b}]^{\cdot} = k\Theta + (1 + R)k\Psi$$

- Photon continuity remains the same

$$\dot{\Theta} = -\frac{k}{3}v_{\gamma b} - \dot{\Phi}$$

- Modification of oscillator equation

$$[(1 + R)\dot{\Theta}]^{\cdot} + \frac{1}{3}k^2\Theta = -\frac{1}{3}k^2(1 + R)\Psi - [(1 + R)\dot{\Phi}]^{\cdot}$$

Oscillator: Take Three

- Combine these to form the not-quite-so simple harmonic oscillator equation

$$c_s^2 \frac{d}{d\eta} (c_s^{-2} \dot{\Theta}) + c_s^2 k^2 \Theta = -\frac{k^2}{3} \Psi - c_s^2 \frac{d}{d\eta} (c_s^{-2} \dot{\Phi})$$

where $c_s^2 \equiv \dot{p}_{\gamma b} / \dot{\rho}_{\gamma b}$

$$c_s^2 = \frac{1}{3} \frac{1}{1 + R}$$

- In a CDM dominated expansion $\dot{\Phi} = \dot{\Psi} = 0$ and the adiabatic approximation $\dot{R}/R \ll \omega = kc_s$

$$[\Theta + (1 + R)\Psi](\eta) = [\Theta + (1 + R)\Psi](0) \cos(k s)$$

Baryon Peak Phenomenology

- Photon-baryon ratio enters in **three** ways
- Overall larger **amplitude**:

$$[\Theta + (1 + R)\Psi](0) = \frac{1}{3}(1 + 3R)\Psi(0)$$

- Even-odd peak **modulation** of effective temperature

$$[\Theta + \Psi]_{\text{peaks}} = [\pm(1 + 3R) - 3R] \frac{1}{3} \Psi(0)$$

$$[\Theta + \Psi]_1 - [\Theta + \Psi]_2 = [-6R] \frac{1}{3} \Psi(0)$$

- Shifting of the **sound horizon** down or ℓ_A up

$$\ell_A \propto \sqrt{1 + R}$$

- Actual effects **smaller** since R evolves

Photon Baryon Ratio Evolution

- Oscillator equation has time evolving mass

$$c_s^2 \frac{d}{d\eta} (c_s^{-2} \dot{\Theta}) + c_s^2 k^2 \Theta = 0$$

- Effective mass is $m_{\text{eff}} = 3c_s^{-2} = (1 + R)$
- Adiabatic invariant

$$\frac{E}{\omega} = \frac{1}{2} m_{\text{eff}} \omega A^2 = \frac{1}{2} 3c_s^{-2} k c_s A^2 \propto A^2 (1 + R)^{1/2} = \text{const.}$$

- Amplitude of oscillation $A \propto (1 + R)^{-1/4}$ decays adiabatically as the photon-baryon ratio changes

Oscillator: Take Three and a Half

- The not-quite-so simple harmonic oscillator equation is a forced harmonic oscillator

$$c_s^2 \frac{d}{d\eta} (c_s^{-2} \dot{\Theta}) + c_s^2 k^2 \Theta = -\frac{k^2}{3} \Psi - c_s^2 \frac{d}{d\eta} (c_s^{-2} \Phi)$$

changes in the gravitational potentials alter the form of the acoustic oscillations

- If the forcing term has a temporal structure that is related to the frequency of the oscillation, this becomes a driven harmonic oscillator
- Term involving Ψ is the ordinary gravitational force
- Term involving Φ involves the $\dot{\Phi}$ term in the continuity equation as a (curvature) perturbation to the scale factor

Potential Decay

- Matter-to-radiation ratio

$$\frac{\rho_m}{\rho_r} \approx 24 \Omega_m h^2 \left(\frac{a}{10^{-3}} \right)$$

of order **unity** at recombination in a low Ω_m universe

- Radiation is not stress free and so **impedes** the growth of structure

$$k^2 \Phi = 4\pi G a^2 \rho_r \Delta_r$$

$\Delta_r \sim 4\Theta$ **oscillates** around a constant value, $\rho_r \propto a^{-4}$ so the Newtonian **curvature decays**.

- General rule: potential decays if the dominant energy component has substantial stress fluctuations, i.e. below the generalized sound horizon or Jeans scale

Radiation Driving

- Decay is timed precisely to **drive** the oscillator - close to fully **coherent**

$$\begin{aligned} [\Theta + \Psi](\eta) &= [\Theta + \Psi](0) + \Delta\Psi - \Delta\Phi \\ &= \frac{1}{3}\Psi(0) - 2\Psi(0) = -\frac{5}{3}\Psi(0) \end{aligned}$$

- **5×** the amplitude of the Sachs-Wolfe effect!
- Coherent approximation is **exact** for a photon-baryon fluid but reality is reduced to $\sim 4\times$ because of **neutrino contribution** to radiation
- Actual **initial conditions** are $\Theta + \Psi = \Psi/2$ for radiation domination but comparison to matter dominated SW correct

Damping

- Tight coupling equations assume a perfect fluid: no viscosity, no heat conduction
- Fluid imperfections are related to the mean free path of the photons in the baryons

$$\lambda_C = \dot{\tau}^{-1} \quad \text{where} \quad \dot{\tau} = n_e \sigma_T a$$

is the conformal opacity to Thomson scattering

- Dissipation is related to the diffusion length: random walk approximation

$$\lambda_D = \sqrt{N} \lambda_C = \sqrt{\eta / \lambda_C} \lambda_C = \sqrt{\eta \lambda_C}$$

the geometric mean between the horizon and mean free path

- $\lambda_D / \eta_* \sim \text{few } \%$, so expect the peaks $:> 3$ to be affected by dissipation

Equations of Motion

- Continuity

$$\dot{\Theta} = -\frac{k}{3}v_\gamma - \dot{\Phi}, \quad \dot{\delta}_b = -kv_b - 3\dot{\Phi}$$

where the photon equation remains unchanged and the baryons follow number conservation with $\rho_b = m_b n_b$

- Euler

$$\begin{aligned}\dot{v}_\gamma &= k(\Theta + \Psi) - \frac{k}{6}\pi_\gamma - \dot{\tau}(v_\gamma - v_b) \\ \dot{v}_b &= -\frac{\dot{a}}{a}v_b + k\Psi + \dot{\tau}(v_\gamma - v_b)/R\end{aligned}$$

where the photons gain an anisotropic stress term π_γ from radiation viscosity and a momentum exchange term with the baryons and are compensated by the opposite term in the baryon Euler equation

Viscosity

- **Viscosity** is generated from radiation **streaming** from hot to cold regions
- Expect

$$\pi_\gamma \sim v_\gamma \frac{k}{\dot{\tau}}$$

generated by streaming, suppressed by **scattering** in a wavelength of the fluctuation. **Radiative transfer** says

$$\pi_\gamma \approx 2A_v v_\gamma \frac{k}{\dot{\tau}}$$

where $A_v = 16/15$

$$\dot{v}_\gamma = k(\Theta + \Psi) - \frac{k}{3}A_v \frac{k}{\dot{\tau}} v_\gamma$$

Oscillator: Penultimate Take

- Adiabatic approximation ($\omega \gg \dot{a}/a$)

$$\dot{\Theta} \approx -\frac{k}{3}v_\gamma$$

- Oscillator equation contains a $\dot{\Theta}$ damping term

$$c_s^2 \frac{d}{d\eta} (c_s^{-2} \dot{\Theta}) + \frac{k^2 c_s^2}{\dot{\tau}} A_v \dot{\Theta} + k^2 c_s^2 \Theta = -\frac{k^2}{3} \Psi - c_s^2 \frac{d}{d\eta} (c_s^{-2} \dot{\Phi})$$

- Heat conduction term similar in that it is proportional to v_γ and is suppressed by scattering $k/\dot{\tau}$. Expansion of Euler equations to leading order in $k\dot{\tau}$ gives

$$A_h = \frac{R^2}{1 + R}$$

since the effects are only significant if the baryons are dynamically important

Oscillator: Final Take

- Final oscillator equation

$$c_s^2 \frac{d}{d\eta} (c_s^{-2} \dot{\Theta}) + \frac{k^2 c_s^2}{\dot{\tau}} [A_v + A_h] \dot{\Theta} + k^2 c_s^2 \Theta = -\frac{k^2}{3} \Psi - c_s^2 \frac{d}{d\eta} (c_s^{-2} \dot{\Phi})$$

- Solve in the adiabatic approximation

$$\Theta \propto \exp(i \int \omega d\eta)$$

$$-\omega^2 + \frac{k^2 c_s^2}{\dot{\tau}} (A_v + A_h) i\omega + k^2 c_s^2 = 0 \quad (1)$$

Dispersion Relation

- Solve

$$\begin{aligned}\omega^2 &= k^2 c_s^2 \left[1 + i \frac{\omega}{\dot{\tau}} (A_v + A_h) \right] \\ \omega &= \pm k c_s \left[1 + \frac{i}{2} \frac{\omega}{\dot{\tau}} (A_v + A_h) \right] \\ &= \pm k c_s \left[1 \pm \frac{i}{2} \frac{k c_s}{\dot{\tau}} (A_v + A_h) \right]\end{aligned}$$

- Exponentiate

$$\begin{aligned}\exp(i \int \omega d\eta) &= e^{\pm i k s} \exp\left[-k^2 \int d\eta \frac{1}{2} \frac{c_s^2}{\dot{\tau}} (A_v + A_h)\right] \\ &= e^{\pm i k s} \exp\left[-(k/k_D)^2\right]\end{aligned}\tag{2}$$

- Damping is exponential under the scale k_D

Diffusion Scale

- Diffusion wavenumber

$$k_D^{-2} = \int d\eta \frac{1}{\dot{\tau}} \frac{1}{6(1+R)} \left(\frac{16}{15} + \frac{R^2}{(1+R)} \right)$$

- Limiting forms

$$\lim_{R \rightarrow 0} k_D^{-2} = \frac{1}{6} \frac{16}{15} \int d\eta \frac{1}{\dot{\tau}}$$

$$\lim_{R \rightarrow \infty} k_D^{-2} = \frac{1}{6} \int d\eta \frac{1}{\dot{\tau}}$$

- Geometric mean between horizon and mean free path as expected from a random walk

$$\lambda_D = \frac{2\pi}{k_D} \sim \frac{2\pi}{\sqrt{6}} (\eta \dot{\tau}^{-1})^{1/2}$$

Thomson Scattering

- Polarization state of radiation in direction $\hat{\mathbf{n}}$ described by the intensity matrix $\langle E_i(\hat{\mathbf{n}})E_j^*(\hat{\mathbf{n}}) \rangle$, where \mathbf{E} is the electric field vector and the brackets denote time averaging.
- Differential cross section

$$\frac{d\sigma}{d\Omega} = \frac{3}{8\pi} |\hat{\mathbf{E}}' \cdot \hat{\mathbf{E}}|^2 \sigma_T ,$$

where $\sigma_T = 8\pi\alpha^2/3m_e$ is the Thomson cross section, $\hat{\mathbf{E}}'$ and $\hat{\mathbf{E}}$ denote the incoming and outgoing directions of the electric field or polarization vector.

- Summed over angle and incoming polarization

$$\sum_{i=1,2} \int d\hat{\mathbf{n}}' \frac{d\sigma}{d\Omega} = \sigma_T$$

Polarization Generation

- Heuristic: incoming radiation shakes an electron in direction of electric field vector $\hat{\mathbf{E}}'$
- Radiates photon with polarization also in direction $\hat{\mathbf{E}}'$
- But photon cannot be longitudinally polarized so that scattering into 90° can only pass one polarization
- Linearly polarized radiation like polarization by reflection
- Unlike reflection of sunlight, incoming radiation is nearly isotropic
- Missing linear polarization supplied by scattering from direction orthogonal to original incoming direction
- Only quadrupole anisotropy generates polarization by Thomson scattering

Acoustic Polarization

- Break down of tight-coupling leads to quadrupole anisotropy of

$$\pi_\gamma \approx \frac{k}{\dot{\tau}} v_\gamma$$

- Scaling $k_D = (\dot{\tau}/\eta_*)^{1/2} \rightarrow \dot{\tau} = k_D^2 \eta_*$
- Know: $k_D s_* \approx k_D \eta_* \approx 10$
- So:

$$\pi_\gamma \approx \frac{k}{k_D} \frac{1}{10} v_\gamma$$

$$\Delta_P \approx \frac{\ell}{\ell_D} \frac{1}{10} \Delta_T$$

Acoustic Polarization

- Gradient of velocity is along direction of wavevector, so polarization is pure E -mode
- Velocity is 90° out of phase with temperature – turning points of oscillator are zero points of velocity:

$$\Theta + \Psi \propto \cos(ks); \quad v_\gamma \propto \sin(ks)$$

- Polarization peaks are at troughs of temperature power

Cross Correlation

- Cross correlation of temperature and polarization

$$(\Theta + \Psi)(v_\gamma) \propto \cos(ks) \sin(ks) \propto \sin(2ks)$$

- Oscillation at twice the frequency
- Correlation: radial or tangential around hot spots
- Partial correlation: easier to measure if polarization data is noisy, harder to measure if polarization data is high S/N or if bands do not resolve oscillations
- Good check for systematics and foregrounds
- Comparison of temperature and polarization is proof against features in initial conditions mimicking acoustic features

CMB Normalization

- Normalization of potential, hence inflationary power spectrum, set by CMB observations, aka COBE or WMAP/Planck normalization
- Angular power spectrum:

$$C_\ell = \frac{4\pi\Delta_T^2(\ell/D_*)}{2\ell(\ell+1)} = \frac{2\pi}{\ell(\ell+1)}\Delta_T^2(\ell/D_*)$$

- $\ell(\ell+1)C_\ell/2\pi = \Delta_T^2$ is commonly used log power
- Sachs-Wolfe effect says $\Delta_T^2 = \Delta_\Phi^2/9$, $\Phi = \frac{3}{5}\mathcal{R}$ initial
- Observed number at recombination

$$\Delta_T^2 = \left(\frac{28\mu\text{K}}{2.725 \times 10^6 \mu\text{K}} \right)^2$$

$$\Delta_\Phi^2 \approx (3 \times 10^{-5})^2$$

$$\Delta_{\mathcal{R}}^2 \approx (5 \times 10^{-5})^2$$

COBE to WMAP to Planck

- Given that the temperature response to an inflationary initial perturbation is known for all k through the Boltzmann solution of the acoustic physics, one can translate Δ_T^2 to $\Delta_{\mathcal{R}}^2$ at the best measured $k \approx \ell/D_*$.
- The CMB normalization was first extracted from COBE at $\ell \sim 10$ or $k \sim H_0$. A low ℓ normalization point suffers from cosmic variance: only $2\ell + 1$ samples of a given ℓ mode.
- WMAP measures very precisely the first acoustic peak at $\ell \approx 200$ which corresponds to $k \sim 0.02 \text{ Mpc}^{-1}$).
- Planck measures out to $\ell = 1000 - 2000$ precisely which corresponds to $k \sim 0.08 \text{ Mpc}^{-1}$
- Taking out the CMB transfer function $\Delta_{\mathcal{R}}^2$ nearly scale invariant with a small red tilt of $n_s - 1 \approx 0.03$

Transfer Function

- **Transfer function** transfers the initial Newtonian curvature to its value today (linear response theory)

$$T(k) = \frac{\Phi(k, a = 1)}{\Phi(k, a_{\text{init}})} \frac{\Phi(k_{\text{norm}}, a_{\text{init}})}{\Phi(k_{\text{norm}}, a = 1)}$$

- Conservation of Bardeen curvature: Newtonian curvature is a **constant** when **stress perturbations are negligible**: above the horizon during radiation and dark energy domination, on all scales during matter domination
- When stress fluctuations dominate, perturbations are stabilized by the **Jeans mechanism**
- Hybrid **Poisson equation**: Newtonian curvature, comoving density perturbation $\Delta \equiv (\delta\rho/\rho)_{\text{com}}$ implies Φ decays

$$(k^2 - 3K)\Phi = 4\pi G a^2 \rho \Delta \sim \eta^{-2} \Delta$$

Transfer Function

- Freezing of Δ stops at η_{eq}

$$\Phi \sim (k\eta_{\text{eq}})^{-2} \Delta_H \sim (k\eta_{\text{eq}})^{-2} \Phi_{\text{init}}$$

- Transfer function has a k^{-2} fall-off beyond $k_{\text{eq}} \sim \eta_{\text{eq}}^{-1}$

$$\eta_{\text{eq}} = 15.7(\Omega_m h^2)^{-1} \left(\frac{T}{2.7K} \right)^2 \text{Mpc}$$

- Small correction since growth with a smooth radiation component is logarithmic not frozen
- Transfer function is a direct output of an Einstein-Boltzmann code

Fitting Function

- Alternately accurate fitting formula exist, e.g. pure CDM form:

$$T(k(q)) = \frac{L(q)}{L(q) + C(q)q^2}$$

$$L(q) = \ln(e + 1.84q)$$


$$C(q) = 14.4 + \frac{325}{1 + 60.5q^{1.11}}$$

$$q = k/\Omega_m h^2 \text{Mpc}^{-1} (T_{\text{CMB}}/2.7K)^2$$

- In $h \text{ Mpc}^{-1}$, the critical scale depends on $\Gamma \equiv \Omega_m h$ also known as the shape parameter

Transfer Function

- Numerical calculation



Baryon Acoustic Oscillations

- Baryons caught up in the acoustic oscillations of the CMB and impart acoustic oscillations to the transfer function. Density enhancements are produced kinematically through the continuity equation $\delta_b \sim (k\eta)v_b$ and hence are out of phase with CMB temperature peaks
- Dissipation of the acoustic oscillations eliminates both the CMB and baryon perturbations – known as Silk damping for the baryons. This suppression and the general fact that baryons are caught up with photons was one of the main arguments for CDM
- Detected first (so far only) in the SDSS LRG survey.
- An excellent standard ruler for angular diameter distance $D_A(z)$ since it does not evolve in redshift in linear theory
- Radial extent of BAO gives $H(z)$ (not yet seen in data)

Massive Neutrinos

- Neutrino dark matter suffers similar effects and hence cannot be the main component of dark matter in the universe
- Relativistic stresses of a light neutrino slow the growth of structure
- Neutrino species with cosmological abundance contribute to matter as $\Omega_\nu h^2 = \sum m_\nu / 94\text{eV}$, suppressing power as $\Delta P/P \approx -8\Omega_\nu/\Omega_m$
- Current data Planck and large scale structure indicate $\sum m_\nu < 0.23\text{eV}$ (95% CL)

Growth Function

- Same physics applies to the dark energy dominated universe
- Under the dark energy sound horizon or Jeans scale, dark energy density frozen. Potential decays at the same rate for all scales

$$G(a) = \frac{\Phi(k_{\text{norm}}, a)}{\Phi(k_{\text{norm}}, a_{\text{init}})} \quad , \quad ' \equiv \frac{d}{d \ln a}$$

- Continuity + Euler + Poisson

$$G'' + \left(1 - \frac{\rho''}{\rho'} + \frac{1}{2} \frac{\rho'_c}{\rho_c}\right) G' + \left(\frac{1}{2} \frac{\rho'_c + \rho'}{\rho_c} - \frac{\rho''}{\rho'}\right) G = 0$$

where ρ is the Jeans unstable matter and ρ_c is the critical density

Dark Energy Growth Suppression

- Pressure growth suppression: $\delta \equiv \delta\rho_m/\rho_m \propto aG$

$$\frac{d^2 G}{d \ln a^2} + \left[\frac{5}{2} - \frac{3}{2} w(z) \Omega_{DE}(z) \right] \frac{dG}{d \ln a} + \frac{3}{2} [1 - w(z)] \Omega_{DE}(z) G = 0,$$

where $w \equiv p_{DE}/\rho_{DE}$ and $\Omega_{DE} \equiv \rho_{DE}/(\rho_m + \rho_{DE})$ with initial conditions $G = 1, dG/d \ln a = 0$

- As $\Omega_{DE} \rightarrow 0$ $g = \text{const.}$ is a solution. The other solution is the decaying mode, eliminated by initial conditions
- As $\Omega_{DE} \rightarrow 1$ $g \propto a^{-1}$ is a solution. Corresponds to a frozen density field.

Power Spectrum Normalization

- Present (or matter dominated) vs inflationary initial conditions (normalized by CMB):

$$\Delta_{\Phi}^2(k, a) \approx \frac{9}{25} \Delta_{\mathcal{R}_i}^2(k_{\text{norm}}) G^2(a) T^2(k) \left(\frac{k}{k_{\text{norm}}} \right)^{n-1}$$

- Density field

$$k^2 \Phi = 4\pi G a^2 \Delta \rho_m$$

$$= \frac{3}{2} H_0^2 \Omega_m \frac{\Delta \rho_m}{\rho_m} \frac{1}{a}$$

$$\Delta_{\Phi}^2 = \frac{9}{4} \left(\frac{H_0}{k} \right)^4 \Omega_m^2 a^{-2} \Delta_m^2$$

$$\Delta_m^2 = \frac{4}{25} \Delta_{\mathcal{R}_i}^2(k_{\text{norm}}) \Omega_m^{-2} a^2 G^2(a) T^2(k) \left(\frac{k}{k_{\text{norm}}} \right)^{n-1} \left(\frac{k}{H_0} \right)^4$$

Antiquated Normalization Conventions


- Current density field on the horizon scale $k = H_0$

$$\delta_H^2 = \frac{4}{25} \Delta_{\mathcal{R}_i}^2(k_{\text{norm}}) \Omega_m^{-2} a^2 G^2(a) = (2G(1)/\Omega_m \times 10^{-5})$$

- σ_8 , RMS of density field filtered by tophat of $8h^{-1}\text{Mpc}$

Power Spectrum

- SDSS data



- Power spectrum defines large scale structure observables: galaxy clustering, velocity field, $\text{Ly}\alpha$ forest clustering, cosmic shear

Velocity field

- Continuity gives the velocity from the density field as

$$\begin{aligned} v &= -\dot{\Delta}/k = -\frac{aH}{k} \frac{d\Delta}{d \ln a} \\ &= -\frac{aH}{k} \Delta \frac{d \ln(aG)}{d \ln a} \end{aligned}$$

- In a Λ CDM model or open model $d \ln(aG)/d \ln a \approx \Omega_m^{0.6}$
- Measuring both the density field and the velocity field (through distance determination and redshift) allows a measurement of Ω_m
- Practically one measures $\beta = \Omega_m^{0.6}/b$ where b is a bias factor for the tracer of the density field, i.e. with galaxy numbers $\delta n/n = b\Delta$
- Can also measure this factor from the redshift space power spectrum - the Kaiser effect where clustering in the radial direction is apparently enhanced by gravitational infall

Redshift Space Power Spectrum

- Kaiser effect is separable from the real space clustering if one measures modes parallel and transverse to the line of sight.
Redshift space distortions only modify the former
- 2D power spectrum in “s” or redshift space

$$P_s(k_{\perp}, k_{\parallel}) = \left[1 + \beta \left(\frac{k_{\parallel}}{k} \right)^2 \right]^2 b^2 P(k)$$

where $k^2 = k_{\parallel}^2 + k_{\perp}^2$ and k_{\perp} is a 2D vector transverse to the line of sight

Power Spectrum Errors

- The precision with which the power spectrum can be measured is ultimately limited by sample variance from having a finite survey volume $V = L^3$. This is basically a mode counting argument. The errors on the power spectrum are given by

$$\left(\frac{\Delta P_s}{P_s} \right)^2 = \frac{2}{N_k}$$

where N_k is the number of modes in a range of $\Delta k_\perp, \Delta k_\parallel$. This is determined by the k -space volume and the fundamental mode of the box $k_0 = 2\pi/L$ which sets the cell size in the volume

$$\left(\frac{\Delta P_s}{P_s} \right)^2 = \frac{2}{\frac{V}{(2\pi)^3} 2\pi k_\perp \Delta k_\perp \Delta k_\parallel}$$

Lyman- α Forest

- QSO spectra absorbed by neutral hydrogen through the Ly α transition.
- The optical depth to absorption is (with ds in physical scale)

$$\tau(\nu) = \int ds x_{\text{HI}} n_b \sigma_{\alpha} \sim \int ds x_{\text{HI}} n_b \Gamma \phi(\nu) \lambda^2$$

where x_{HI} is the neutral fraction, $\Gamma = 6.25 \times 10^8 \text{ s}^{-1}$ is the transition rate and $\lambda = 1216 \text{ \AA}$ is the Ly α wavelength and $\phi(\nu)$ is the Lorentz profile. For radiation at a given emitted frequency ν_0 above the transition, it will redshift through the transition

- Resonant transition: lack of complete absorption, known as the lack of a Gunn-Peterson trough indicates that the universe is nearly fully ionized $x_{\text{HI}} \ll 1$ out to the highest redshift quasar $z \sim 6$; indications that this is near the end of the reionization epoch

Lyman- α Forest

- In ionization equilibrium, the Gunn-Peterson optical depth is a tracer of the underlying baryon density which itself is a tracer of the dark matter $\tau_{GP} \propto \rho_b^2 T^{-0.7}$ with $T(\rho_b)$.

$$\frac{d(1 - x_{\text{HI}})}{dt} = -x_{\text{HI}} \int d\nu \frac{4\pi J_\nu}{h\nu} \sigma_\nu + (1 - x_{\text{HI}})^2 n_b R$$

where σ_ν is the photoionization cross section (sharp edge at threshold and falling in frequency means $J_\nu \approx J_{21}$) and $R \propto T^{-0.7}$ is the recombination coefficient.

- Given an equation of state from simulations of $p \propto \rho^\gamma$

$$x_{\text{HI}} \propto \frac{\rho_b R}{J_{21}} \propto \frac{\rho_b T^{-0.7}}{J_{21}}, \quad \tau_{GP} \propto \frac{\rho_b^{2-0.7(\gamma-1)}}{J_{\text{HI}}}$$

- Clustering in the Ly α forest reflects the underlying power spectrum modulo an overall ionization intensity J_{21}

Gravitational Lensing

- Gravitational potentials along the line of sight $\hat{\mathbf{n}}$ to some source at comoving distance D_s lens the images according to (flat universe)

$$\phi(\hat{\mathbf{n}}) = 2 \int dD \frac{D_s - D}{DD_s} \Phi(D\hat{\mathbf{n}}, \eta(D))$$

remapping image positions as

$$\hat{\mathbf{n}}^I = \hat{\mathbf{n}}^S + \nabla_{\hat{\mathbf{n}}} \phi(\hat{\mathbf{n}})$$

- Since absolute source position is unknown, use image distortion defined by the Jacobian matrix

$$\frac{\partial n_i^I}{\partial n_j^S} = \delta_{ij} + \psi_{ij}$$

Weak Lensing

- Small image distortions described by the convergence κ and shear components (γ_1, γ_2)

$$\psi_{ij} = \begin{pmatrix} \kappa - \gamma_1 & -\gamma_2 \\ -\gamma_2 & \kappa + \gamma_1 \end{pmatrix}$$

where $\nabla_{\hat{\mathbf{n}}} = D\nabla$ and

$$\psi_{ij} = 2 \int dD \frac{D(D_s - D)}{D_s} \nabla_i \nabla_j \Phi(D\hat{\mathbf{n}}, \eta(D))$$

- In particular, through the Poisson equation the convergence (measured from shear) is simply the projected mass

$$\kappa = \frac{3}{2} \Omega_m H_0^2 \int dD \frac{D(D_s - D)}{D_s} \frac{\Delta(D\hat{\mathbf{n}}, \eta(D))}{a}$$