

Evolution of a k -mode

Take the initial conditions for a given k -mode from CMB PS# 3 and evolve the equations of motion in linear perturbation theory for a fully ionized universe.

In the notation of the previous problem sets and class notes, your fundamental variables are for the density perturbations

$$\Delta_\gamma, \Delta_b, \Delta_c \quad (1)$$

for the fluid velocities

$$v_\gamma, v_b, v_c, \quad (2)$$

[if you care, these are all in comoving gauge; although v_b is not technically needed for the approximation you are asked to solve, keep it anyway so that you can generalize your code later]. Your auxiliary variables are: the Newtonian curvature Φ , the Newtonian potential Ψ , the conformal time derivative of the Bardeen curvature $\dot{\zeta}$, the Newtonian temperature perturbation Θ , the anisotropic stress of the photons π_γ and the entropy perturbation in the photon-baryon system σ .

Explicitly, the set of coupled linear differential equations are:

(1) Continuity

$$\begin{aligned} \dot{\Delta}_\gamma &= -\frac{4}{3}(kv_\gamma + 3\dot{\zeta}) \\ \dot{\Delta}_b &= -(kv_b + 3\dot{\zeta}) \\ \dot{\Delta}_c &= -(kv_c + 3\dot{\zeta}) \end{aligned} \quad (3)$$

(2) Euler

$$\begin{aligned} \dot{v}_\gamma &= -\frac{R}{1+R} \frac{\dot{a}}{a} v_\gamma + \frac{1}{1+R} k\Theta + k\Psi - \frac{1}{3} \frac{R}{(1+R)^2} k\sigma - \frac{1}{6} \frac{1}{(1+R)} k\pi_\gamma \\ \dot{v}_b &= \dot{v}_\gamma \\ \dot{v}_c &= -\frac{\dot{a}}{a} v_c + k\Psi \end{aligned} \quad (4)$$

For which you will need the definitions of the auxiliary parameters:

$$k^2\Phi = 4\pi G a^2 \sum_i \Delta_i \rho_i \quad (5)$$

$$k^2(\Psi + \Phi) = -\frac{8}{3} \pi G a^2 \rho_\gamma \pi_\gamma \quad (6)$$

$$\Theta = \frac{1}{4} \Delta_\gamma - \frac{\dot{a}}{a} v/k \quad (7)$$

$$v = \frac{\sum_i (\rho_i + p_i) v_i}{\sum_i (\rho_i + p_i)} \quad (8)$$

$$\dot{\zeta} \left(\frac{\dot{a}}{a} \right)^{-1} = -\frac{w}{(1+w)} \left(\Delta_\gamma - \frac{2}{3} \pi_\gamma \right) \quad (9)$$

$$w = \frac{1}{3} \frac{\rho_\gamma}{\rho} \quad (10)$$

$$\sigma = (k\dot{\tau}^{-1}) R v_\gamma \quad (11)$$

$$\pi_\gamma = \frac{32}{15} (k\dot{\tau}^{-1}) v_\gamma. \quad (12)$$

where the sums are over the three particle species. You can and should verify all these relationships yourselves from your PS's and notes since I am prone to make sign errors, etc!

Test your code.

Artificially set $\sigma = \pi_\gamma = 0$ to make the system dissipationless.

- Choose a $k \gg \eta_{\text{eq}}^{-1}$ and plot the evolution of the fundamental and auxiliary parameters. In particular what is the amplitude of the acoustic oscillation in Θ in terms of $\zeta(0)$? check the answer with the solution given in class. What is the behavior of the Newtonian potential Φ ? again check this against what we learned in class.

- Choose a $k \ll \eta_{\text{eq}}^{-1}$ and follow the evolution well past horizon crossing (ignoring recombination). What is the value of Φ and Ψ compared with $\zeta(0)$, check your answer. How does the amplitude of the oscillation in $\Theta + \Psi$ behave as you cross $R = 1$, verify the qualitative behavior discussed in class.

Turn dissipation back on.

- For the $k \gg \eta_{\text{eq}}^{-1}$ case, when does the acoustic oscillation dissipate, compare that with the random walk (or full dissipation) calculation in class.